

Omeprazole, Acid Reducer

Procure Omeprazole delayed-release tablets belong to the class of Proton Pump Inhibitors (PPIs). These tablets are commonly used to reduce the production of stomach acid and treat conditions such as gastroesophageal reflux disease (GERD), stomach ulcers, and excessive stomach acid secretion.



Benefits:

- Treats Frequent Heartburn[†]
- Acid Reducer[†]
- 20mg Delayed-Release Tablets
- Three 14-Day Courses Treatment

[†] These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Visit Us Online at:
www.ProCureProducts.com

Product	Strength	Reorder #	Units
Omeprazole	20mg	PCOTC120	42 Pills/Box, 24 Bottles/Case

* Compare to active ingredient in Prilosec OTC

PROCURE PRODUCTS

Dear Valued TwinMed Customer,

TwinMed is proud to announce the addition of Omeprazole OTC 20mg Tablets to the Procure line of products.

The Procure Omeprazole 20mg is equivalent to the National Brand Prilosec OTC and they may be used interchangeably.

We wanted to ensure any general questions about the Procure brand are addressed.

Announcing a
NEW Product



Treats
Frequent
Heartburn!

Q. How much active ingredient of Omeprazole is in the Procure Omeprazole and Prilosec Brand?

A. Both Procure Brand and Prilosec OTC brand contain **20 mg of Omeprazole**. In addition, any previous brands of Omeprazole you were supplied by TwinMed also contained 20mg of Omeprazole.

Q. Is the Procure Brand Omeprazole Tablet crushable? The Procure brand is a tablet just like the Prilosec brand.

A. All Omeprazole OTC products on the market today including capsules states on the label, to swallow whole or **“DO NOT CRUSH”**. Questions about crushing tablets should be reviewed and consulted with a Healthcare Professional.

PROCURE
HEALTHCARE PRODUCTS

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole USP 20 mg</p> <p>Purpose Acid reducer</p> <p>Use</p> <ul style="list-style-type: none"> • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you have:</p> <ul style="list-style-type: none"> • trouble or pain swallowing food, vomiting with blood, or bloody or black stools • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or • frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> • had heartburn over 3 months. This may be a sign of a more serious condition. • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain 	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months • you get diarrhea • you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> • for adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> • swallow 1 tablet with a glass of water before eating in the morning • take every day for 14 days • do not take more than 1 tablet a day • do not use for more than 14 days unless directed by your doctor <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> • you may repeat a 14-day course every 4 months 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • do not take for more than 14 days or more often than every 4 months unless directed by a doctor • children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. <p>Other information</p> <ul style="list-style-type: none"> • read the directions and warnings before use • keep the carton. It contains important information. • store at 20 to 25° C (68 to 77° F) and protect from moisture <p>Inactive ingredients anhydrous lactose, hydroxymethylcellulose acetate succinate, iron oxide red, iron oxide yellow, lactose monohydrate, methylcellulose, monoethanolamine, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, sodium stearate, sodium stearyl fumarate, talc, triethyl citrate and titanium dioxide.</p> <p>The imprinting ink contains ammonium hydroxide, black iron oxide, n-butyl alcohol, propylene glycol and shellac.</p> <p>Questions or Comments? Call toll free 1-800-818-4535 weekdays.</p> <p>Safety Feature - Do not use if printing seal under cap is broken or missing.</p>
---	--	--

Compare to
the active
ingredient of
Prilosec OTC®

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium). Acid reducer</p> <p>Purpose</p> <ul style="list-style-type: none"> • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Use</p> <ul style="list-style-type: none"> • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> • trouble or pain swallowing food, vomiting with blood, or bloody or black stools • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> • had heartburn over 3 months. • This may be a sign of a more serious condition. • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> • your heartburn continues or worsens • you need to take this product for more than 14 days 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> • you need to take more than 1 course of treatment every 4 months • you get diarrhea • you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> • for adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> • swallow 1 tablet with a glass of water before eating in the morning • take every day for 14 days • do not take more than 1 tablet a day • do not use for more than 14 days unless directed by your doctor <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> • you may repeat a 14-day course every 4 months • do not take for more than 14 days or more often than every 4 months unless directed by a doctor <ul style="list-style-type: none"> • children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. <p>Other information</p> <ul style="list-style-type: none"> • read the directions and warnings before use • keep the carton. It contains important information. • store at 20-25° C (68-77° F) and protect from moisture
---	--



Q. The Procure brand only states Omeprazole 20mg on the label and the Prilosec brand states “Equivalent to 20.6mg Omeprazole Magnesium” – is there a difference?

A. No. Both the Prilosec Brand and Procure Brand contain 20.6mg of the compound Omeprazole Magnesium. Of this total, each tablet contains 20 mg of Omeprazole and 0.6 mg of Magnesium. However, due to IP (intellectual Property rights) and FDA labeling guidelines for generic Prilosec tablets, we are only allowed to list the Omeprazole 20mg in the drug facts and not the 0.6mg of Magnesium.

Q. If there is 0.6mg of Magnesium, why is it not listed in the ingredients?

A. The 0.6mg is considered insignificant by FDA standards. To put the amount of Magnesium in perspective, the Recommended Daily Allowance (RDA) for Magnesium in adults is 270 mg – 400 mg for males and 280 mg – 300 mg for females. The FDA does not require an OTC drug label to state the Magnesium content for products containing less than 8 mg of Magnesium because less than 8 mg the FDA considers insignificant and risk free.